

# IOWA STATE UNIVERSITY

## Digital Repository

---

Integrated Crop Management News

Agriculture and Natural Resources

---

4-16-2007

## Black cutworms and the hard freeze

Marlin E. Rice

*Iowa State University*, [merice@iastate.edu](mailto:merice@iastate.edu)

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Entomology Commons](#), and the [Meteorology Commons](#)

---

### Recommended Citation

Rice, Marlin E., "Black cutworms and the hard freeze" (2007). *Integrated Crop Management News*. Paper 1112.  
<http://lib.dr.iastate.edu/cropnews/1112>

This Article is brought to you for free and open access by the Agriculture and Natural Resources at Digital Repository @ Iowa State University. It has been accepted for inclusion in Integrated Crop Management News by an authorized administrator of Digital Repository @ Iowa State University. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

# INTEGRATED CROP MANAGEMENT

  

Get the latest research-based information on crops. [Sign up to be notified](#) when new content is available!

ICM > 2007 > IC-498 (6) -- April 16, 2007

## Current Newsletter

You are viewing **archives** for the newsletter from 1993-2007. For current news, see [Integrated Crop Management News](#).

## Archives 1993-2007



Announcements



Crop Production



Insects and Mites



Pesticide Education



Plant Diseases



Soils



Weed Management

[Image Gallery](#)

## Printable Version

Printable version of this page

## Related Articles

Black cutworm 2007 cutting dates predicted  
**May 7, 2007**

Early-season insects in your corn field  
**May 15, 2006**

Black cutworms return, 2006 cutting dates predicted  
**May 1, 2006**

Seed treatments, black

## Black cutworms and the hard freeze

*by Marlin E. Rice, Department of Entomology*

Each spring, black cutworm moths migrate into Iowa from southern states, and they lay eggs in soybean stubble (a preferred site) and corn stubble. Also, at the same time, we invariably experience some early spring cold temperatures that drop below freezing. When these two events occur simultaneously, it doesn't take long before people begin asking what the freezing temperatures did to the black cutworms.

It is fairly well known that many insects that migrate into the northern states are fairly adept at finding protected shelter to ride out the cold temperatures. But what about any black cutworm eggs that are laid in a field? Wouldn't they be exposed to the cold and possibly be killed? Well, the answer to the first question is yes, they would be exposed to the cold, but I am not aware of an answer to the second question. Therefore, I set up a quick experiment to determine what effect a natural freeze would have on black cutworm eggs.

On April 4, I obtained laboratory-laid black cutworm eggs from the USDA Corn Insects Lab at Iowa State University. These eggs were two days old and laid on paper toweling. Twenty-four clusters of eggs (60-150 each) were cut from the paper toweling and then glued to manila envelopes. Twelve, or half, of the envelopes were kept indoors at 72 °F while the other 12 were placed outside overnight in a soybean field.



©MARLIN E. RICE

cutworms, and white grubs: What should you expect?

**April 24, 2006**

Black cutworms, dingy cutworms, and predicted cutting dates

**May 16, 2005**

Seed treatments may not stop cutworms

**May 9, 2005**

First cutting dates predicted for black cutworm

**May 17, 2004**

Blacks and dingys: confusing cutworms

**May 17, 2004**

Preventive insecticides in corn for cutworms

**May 19, 2003**

Black cutworm trapping is underway

**May 12, 2003**

### *Black cutworm eggs. (Marlin Rice)*

The outside temperatures at 6 p.m. were 33 °F; at 10:30 p.m., it had dropped to 28 °F, and at 6:30 a.m. on April 5 it was 19 °F. The outside egg clusters were then brought indoors and kept with the "check" eggs for another five days, at which time the number of hatched eggs were counted in both groups.

So what did I find? The percent of larvae hatching between the two groups was nearly identical; 61.7 percent of eggs kept inside hatched, while 72.3 percent of those kept outside overnight hatched. The eight hours of freezing temperatures, with a low of 19 °F, appeared to have no effect on the survival of the black cutworm eggs and subsequent larval hatch! So it would appear that the black cutworm is well adapted to surviving brief encounters of very cold temperatures. An obvious next question is to determine the lethal minimum temperature, or how cold does it need to get before black cutworm eggs start freezing? This is an answer I don't know.

*Marlin E. Rice is a professor of entomology with extension and research responsibilities in field and forage crops.*

This article originally appeared on page 113 of the IC-498 (6) -- April 16, 2007 issue.

Updated 04/18/2007 - 2:43pm